

## Empathy for the Unvaccinated

Dear Editor:

When Dr Ehlenbach shared the news of his father's diagnosis of advanced lung cancer with colleagues, he was frequently asked by these physicians if his father had been a smoker. This question reveals an implicit desire to separate the complicit from the innocent, to fully inform their reaction to the news. Regardless, it is sad that his father was sick and that he died, full stop. Blaming patients for their health problems is a practice that predates scientific medicine, but it seems to have become worse over time. We invoke patients' failure to control diabetes as an explanation for limb loss, lack of discipline as the etiology of obesity, and "medication noncompliance" as the cause of runaway hypertension. Improving health through behavioral modification does not require vilification of patients. But our professions' bad habits threaten to destroy expressions of empathy required for a therapeutic relationship.

Nowhere is this more seditious or prevalent than for our patients with serious illness from COVID-19. The vast majority of patients with COVID currently filling intensive care units in Wisconsin have not received a COVID vaccine, and most of them would not be severely ill had they done so. Tens of thousands of Americans will die because they did not get vaccinated. For an already exhausted workforce, the notion that we could feel less exhausted, safer, and less burned out if people would simply take a free, safe, and highly effective vaccination is as straightforward as it is maddening that so many have not done so. The light at the end of the tunnel has dimmed. It is hard to watch people die from preventable illness. But this is what we do and have always done. It is challenging to care for people whose illness may harm you, yet this is also part of the job – consider tuberculosis, hepatitis C, and HIV. In return for our care, we receive relatively high pay and high status. We are not qualified to be judge and jury. In making these judgements we are likely to make mistakes.

We are desperately in need of a better narrative. Shifting narratives to support empathy can improve care as it has in substance use disorder. When we moved from considering addiction as a moral failing to conceptualizing it as a disease, it fundamentally altered our approach. Shaming our patients for failure to vaccinate, or routinely expressing anger to colleagues that these patients are sick because they are dumb, simply inflames the toxic divisions that got them to this

place. To start, we need a different target for our well-justified anger. So we should focus our anger on those who misled them. Elected leaders and media personalities who spread misinformation and social media platforms like Facebook that amplify it have led our patients to make choices that are harming them and those around them. Next, we need to reframe our feelings about critically ill adult patients with COVID. It is tragic and it is exhausting, and we should acknowledge this. A better narrative is that we are sad that our patients succumbed to lies spread by people they trusted. We are flummoxed that they trusted them more than they trusted us. We will do better to be trustworthy, but for now we will fight their illness with them.

People will always be sick, humans will always make bad choices, and we will always be here to take care of them. It's time to have more empathy for the unvaccinated.

*Margaret L. Schwarze, MD, MPP; William J. Ehlenbach, MD, MSc*

• • •

**Author Affiliations:** Department of Medicine, University of Wisconsin School of Medicine and Public Health (UW SMPH), Madison, Wisconsin (Ehlenbach, Schwarze); Department of Surgery and Department of Medical History and Bioethics, UW SMPH (Schwarze).

**Corresponding Author:** William J. Ehlenbach, MD, MSc, Assistant Professor, University of Wisconsin School of Medicine and Public Health, 2281 UW Medical Foundation, Rm 5245 Centennial Bldg, 1685 Highland Ave, Madison, WI 53705; email [wjehlen@medicine.wisc.edu](mailto:wjehlen@medicine.wisc.edu).

**Funding/Support:** None declared.

**Financial Disclosures:** None declared.

## Racial and Ethnic Differences in Zoster Vaccine Uptake: A Cross-Sectional Study in a Veterans Health Administration Primary Care Clinic

Dear Editor:

Zoster vaccine uptake has been disappointing (34.5% of the target population) and marred by racial and ethnic disparities.<sup>1,2</sup> However, studies of uptake generally have limitations. Most are survey-based (and subject to self-report bias) and based largely on the discontinued live vaccine

(not the currently available recombinant vaccine).<sup>1,3</sup> Furthermore, much of recent literature describes the situation 3 years ago when the recombinant vaccine was in shortage.<sup>1</sup>

These limitations raise a question: Do these racial and ethnic disparities persist? Insight into that question may be gleaned from a quality improvement project that we initiated to improve zoster vaccine uptake. Our baseline findings overcome those limitations. Our findings are current, record-based, and reflect the recombinant vaccine. Our findings are from a Veterans Health Administration clinic (where insurance and access are not barriers) and may shed light on the question of persistence of disparities, even when those barriers are absent.<sup>4</sup>

We queried the records of the Omaha primary care clinic of the Veterans Health Administration Nebraska-Western Iowa Health Care System for receipt of recombinant zoster vaccine since October 1, 2017. We included patients at least 50 years old on October 1, 2017 (close to the recombinant vaccine approval date) seen in the clinic October 1, 2020-July 5, 2021.

Our population of 10,323 was predominantly male (93.8%); 81.2% were non-Hispanic White, 10.7% were non-Hispanic Black, and 1.5% were Hispanic White. The prevalence of complete vaccination (2 doses) was 39.8% (females 34.7%, males 40.1%). Complete vaccination was 43.3% in non-Hispanic White patients, 33.8% in Hispanic White patients, and 24.9% in non-Hispanic Black patients. Receipt of at least 1 COVID-19 vaccine dose was 80.1%, 78.2%, and 82.2%, respectively.

A 39.8% prevalence of complete vaccination was higher than generally reported for zoster vaccine uptake.<sup>1,2</sup> Conceivably, this could reflect our study population: individuals seen in a clinic with vaccine reminders, standing vaccine orders, on-site vaccine, and no charge for vaccine.<sup>4,5</sup> Racial and ethnic disparities are consistent with most, but not all, of the literature.<sup>1,2</sup>

The contrast between zoster vaccine disparities and their absence with COVID-19 vaccine (for which awareness was extraordinarily high) supports the hypothesis that zoster vaccine disparities arise from disparities in awareness.<sup>2</sup>

Our baseline data confirm the appropriateness of our choice of zoster vaccine uptake as a quality improvement project, showing an opportunity for improving uptake and an opportunity to address factors other than insurance and access that account for racial and ethnic disparities.

*Marvin J. Bittner, MD; Gia Thinh D. Truong, BS; Zachary A. Creech, BS*

*continued on page 171*

One trend that has emerged from the last two years of a pandemic is the increase in attention to rural health and the promotion of telehealth options of care, which may be ideal for people who live long distances away from health care. The US Department of Health and Human Services designated \$1 billion to improve the COVID-19 response in rural areas. The money is designated to increase vaccine use and to support small hospitals by increasing infrastructure.<sup>10</sup> We may not be finished with this pandemic, but hopefully, some progress in improving rural health will remain.

## REFERENCES

1. What is rural? Economic Research Service. US Department of Agriculture. Accessed Sept 2, 2021. <https://www.ers.usda.gov/topics/rural-economy-population/rural-classifications/what-is-rural/>
2. Warshaw R. Health disparities affect millions in rural US Communities. *AAMC News*. October 31, 2017. Accessed Sept 2, 2021. <https://www.aamc.org/news-insights/health-disparities-affect-millions-rural-us-communities>
3. Solenkova N. Op-ed: rural hospitals struggling with COVID care—ICUs stretched past the breaking point. *MedPage Today*. November 30, 2020. Accessed September 2, 2021. <https://www.medpagetoday.com/infectiousdisease/covid19/89930>
4. Gupta S, Hayek SS, Wang W, et al. Factors associated with death in critically ill patients with coronavirus disease 2019 in the US. *JAMA Intern Med*. 2020;180(11):1436–1447. doi:10.1001/jamainternmed.2020.3596
5. Callaghan T, Lueck JA, Trujillo KL, Ferdinand AO. Rural and urban differences in COVID-19 prevention behaviors. *J Rural Health*. 2021;37(2):287–295. doi:10.1111/jrh.12556
6. Schrage S. Rural health in Wisconsin—looking to the future. *WMJ*. 2019;117(5):192–193.
7. Park KY, Eithun BL, Havlena J, et al. Driving time to trauma centers for children living in Wisconsin. *WMJ*. 2021;120(3):174–177.
8. McClure DK, Cullinane DC, Maldonado, IL. Trends in mortality at a level II rural trauma center. *WMJ*. 2021;120(3):178–182.
9. Anderson D, Beinhoff P, Ruffalo L. Rural residence predicts lower cardiac mortality and better mental health outcomes. *WMJ*. 2021;120(3):183–187.
10. Rubin R. Funding for rural COVID-19 care. *JAMA*. 2021;325(24):2426. doi:10.1001/jama.2021.8995

## Letters to the Editor

continued from page 168

### REFERENCES

1. Terlizzi EP, Black LI. Shingles vaccination among adults aged 60 and over: United States, 2018. NCHS Data Brief, no 370. National Center for Health Statistics; 2020. Accessed August 19, 2021. <https://www.cdc.gov/nchs/products/databriefs/db370.htm>
2. Elekwachi O, Wingate LT, Clarke Tasker V, et al. A review of racial and ethnic disparities in immunizations for elderly adults. *J Prim Care Community Health*. 2021;12; 21501327211014071. doi:10.1177/21501327211014071
3. Vogelsang EM, Polonijo AN. Social determinants of shingles vaccination in the United States. *J Gerontol B Psychol Sci Soc Sci*. 2021;gbab074. doi:10.1093/geronb/gbab074
4. Straits-Tröster KA, Kahwati LC, Kinsinger LS, Orelien J, Burdick MB, Yevich SJ. Racial/ethnic differences in influenza vaccination in the Veterans Affairs healthcare system. *Am J Prev Med*. 2006;31(5):375–382. doi:10.1016/j.amepre.2006.07.018

5. Brewer NT, Chapman GB, Rothman AJ, Leask J, Kempe A. Increasing vaccination: putting psychological science into action. *Psychol Sci Public Interest*. 2017;18(3):149–207. doi:10.1177/1529100618760521

...

**Corresponding Author:** Marvin J. Bittner, MD, Veterans Health Administration Nebraska-Western Iowa Healthcare System, 4101 Woolworth Ave (111D), Omaha, NE 68105; email [marvin.bittner@va.gov](mailto:marvin.bittner@va.gov); ORCID ID 0000-0002-2464-8819.

**Author Affiliations:** Veterans Health Administration Nebraska-Western Iowa Healthcare System, Omaha, Nebraska (Bittner); Creighton University School of Medicine, Omaha, Nebraska (Bittner, Truong, Creech).

**Financial Disclosures:** None declared.

**Funding/Support:** None declared.

## Let us hear from you!

If an article strikes a chord or you have something on your mind related to medicine, share it with your colleagues. Email your letter to the editor to [wmj@med.wisc.edu](mailto:wmj@med.wisc.edu).

# WMJ

Let **MatchingDonors.com** Help You  
**FIND YOUR LIVING ORGAN DONOR TODAY**

Search over **15,000+** Willing Donors

Many of our patients receive their transplant in just **6 months**

a 501(c)(3) nonprofit organization  
1-781-821-2204

MatchingDonors



advancing the art & science of medicine in the midwest

**WMJ**

*WMJ* (ISSN 1098-1861) is published through a collaboration between The Medical College of Wisconsin and The University of Wisconsin School of Medicine and Public Health. The mission of *WMJ* is to provide an opportunity to publish original research, case reports, review articles, and essays about current medical and public health issues.

© 2021 Board of Regents of the University of Wisconsin System and The Medical College of Wisconsin, Inc.

**Visit [www.wmjonline.org](http://www.wmjonline.org) to learn more.**